

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP05/002635

International filing date: 04 March 2005 (04.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: GB  
Number: 0405198.3  
Filing date: 08 March 2004 (08.03.2004)

Date of receipt at the International Bureau: 26 April 2005 (26.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



EP05/2635



INVESTOR IN PEOPLE


The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.


Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated 29 December 2004





**The  
Patent  
Office**

PCT/EP200 5/002635  
P01/7700 0.00-0405198.3 NONE

**1/77**

**Request for grant of a patent**

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

08 MAR 2004

**The Patent Office**

Cardiff Road  
Newport  
South Wales  
NP10 8QQ

1. Your Reference **RK/PB60773P**

2. Patent application number **0405198.3**  
(The Patent office will fill in this part)

3. Full name, address and postcode of the or of each applicant (underline all surnames) **GLAXO GROUP LIMITED  
GLAXO WELLCOME HOUSE  
BERKELEY AVENUE  
GREENFORD  
MIDDLESEX  
UB6 ONN  
GB**

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its corporation **GB**

473587003

4 Title of the invention **COMPOUNDS**

5 Name of your agent (if you know one) **RIE KONDO**

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode) **GLAXOSMITHKLINE  
CORPORATE INTELLECTUAL PROPERTY  
CN925.1  
980 GREAT WEST ROAD  
BRENTFORD  
MIDDLESEX  
TW8 9GS, GB**

Patents ADP number (if you know it)

8072555004

6. Priority: Complete this section if you are declaring priority from one or more earlier patent applications, filed in the last 12 months	Country	Priority application number (if you know it)	Date of Filing (day / month / year)

7. Divisionals: etc Complete this section only if this application is a divisional application or resulted from an entitlement dispute (see note f)	Number of earlier application	Date of filing (day / month / year)

8. Is a Patents Form 7/77 (Statement of inventorship and of right to grant of a patent) required in support of this request? **YES**

Answer YES if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body

Otherwise answer NO See note (d)

## Patents Form 1/77

9. Accompanying documents: A patent application must include a description of the invention. Not counting duplicates, please enter the number of pages of each item accompanying this form:

Continuation sheets of this form	-
Description	20
Claim(s)	6
Abstract	-
Drawing(s)	-

10. If you are also filing any of the following, state how many against each item

### Priority Documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patent Form 9/77*)

Request for substantive examination (*Patent Form 10/77*)

Any other documents(*please specify*)

11. I/We request the grant of a patent on the basis of this application

Signature

  
RIE KONDO. Date 8 March 2004  
**AGENT FOR THE APPLICANTS**

12. Name and daytime telephone number and e-mail address, if any, of person to contact in the United Kingdom
- Jean Harney +44 (0)208 047 4420  
jean.l.harney@gsk.com

### Warning

*After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked*

### Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.*
- Write your answers in capital letters using black ink or you may type them.*
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form*
- If you have answered "Yes" in part 8, a Patents Form 7/77 will need to be filed.*
- Once you have filled in the form you must remember to sign and date it.*
- Part 7 should only be completed when a divisional application is being made under section 15(4), or when an application is being made under section 8(3), 12(6) or 37(4) following an entitlement dispute. By completing part 7 you are requesting that this application takes the same filing date as an earlier UK application. If you want the new application to have the same priority date(s) as the earlier UK application, you should also complete part 6 with the priority details.*


## COMPOUNDS

The present invention relates to novel compounds, processes for their preparation, intermediates used in these processes, pharmaceutical compositions containing them and their use in therapy, as modulators of dopamine D<sub>3</sub> receptors, in particular as antipsychotic agents.

WO 2002/40471 (SmithKline Beecham) discloses certain benzodiazepine compounds having activity at the dopamine D<sub>3</sub> receptor.

A new class of compounds which have affinity for dopamine receptors, in particular the dopamine D<sub>3</sub> receptor, has been found. These compounds have potential in the treatment of conditions wherein modulation, especially antagonism/inhibition, of the D<sub>3</sub> receptor is beneficial, e.g. as antipsychotic agents or to treat drug dependency.

The present invention provides a compound of formula (I) or a salt thereof:




wherein

- R<sub>1</sub> and R<sub>4</sub> are independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, C<sub>1-2</sub>alkyl, C<sub>1</sub>alkoxy, haloC<sub>1-2</sub>alkyl, haloC<sub>1</sub>alkoxy, hydroxy, cyano and nitro;
- R<sub>2</sub> and R<sub>3</sub> are independently selected from the group consisting of:  
 halogen, hydroxy, cyano, nitro, C<sub>1-4</sub>alkyl, haloC<sub>1-4</sub>alkyl, C<sub>3-6</sub>cycloalkyl, C<sub>1-4</sub>alkoxy, haloC<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkylthio, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, C<sub>3-6</sub>cycloalkylC<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-4</sub>alkoxycarbonylC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylsulfonyl, C<sub>1-4</sub>alkylsulfonyloxy, haloC<sub>1-4</sub>alkylsulfonyl, haloC<sub>1-4</sub>alkylsulfonyloxy, C<sub>1-4</sub>alkylsulfonylC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylsulfonamido, C<sub>1-4</sub>alkylsulfonamidoC<sub>1-4</sub>alkyl, heterocyclyl, aryl, arylC<sub>1-4</sub>alkoxy, aryloxy, arylthio, arylmethyl, aroyl, aryloxymethyl, arylsulfonyl, aryl-NR' (wherein R' is hydrogen or C<sub>1-4</sub>alkyl), arylsulfonyloxy, arylsulfonylC<sub>1-4</sub>alkyl, arylsulfonamido, arylcarboxamido, arylsulfonamidoC<sub>1-4</sub>alkyl, arylcarboxamidoC<sub>1-4</sub>alkyl, aroylC<sub>1-4</sub>alkyl, arylC<sub>1-4</sub>alkanoyl, a group R<sub>11</sub>CON(R<sub>12</sub>)(CH<sub>2</sub>)<sub>r</sub>, R<sub>11</sub>R<sub>12</sub>NCO(CH<sub>2</sub>)<sub>r</sub> or R<sub>11</sub>R<sub>12</sub>NSO<sub>2</sub>(CH<sub>2</sub>)<sub>r</sub> (in which r is 0, 1, 2, 3 or 4, and each of R<sub>11</sub> and R<sub>12</sub> is independently hydrogen or C<sub>1-4</sub>alkyl, or in the groups R<sub>11</sub>CON(R<sub>12</sub>)(CH<sub>2</sub>)<sub>r</sub>, R<sub>11</sub>R<sub>12</sub>NCO(CH<sub>2</sub>)<sub>r</sub> and

$R_{11}R_{12}NSO_2(CH_2)_t$ ,  $R_{11}CONR_{12}$  or  $R_{11}R_{12}N$  together form a 4-, 5-, 6- or 7-membered azacyclic group optionally containing one additional O, N or S atom in the azacycle and having 3-8 carbon atoms (including the carbon atoms contained in any optional substituent(s) of the azacycle)); wherein in any group containing an aryl moiety, the aryl may be substituted by one, two or three groups selected from the group consisting of halogen, hydroxy, cyano, nitro, amino,  $C_{1-4}$ alkyl, halo $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy, halo $C_{1-4}$ alkoxy,  $C_{1-4}$ alkylenedioxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkylsulfonyl, halo $C_{1-4}$ alkylsulfonyl,  $C_{1-4}$ alkylamino,  $C_{1-4}$ dialkylamino,  $R_{13}R_{14}NCO$  (in which  $R_{13}$  and  $R_{14}$  are independently hydrogen or  $C_{1-4}$ alkyl, or  $R_{13}R_{14}N$  together form a 4-, 5-, 6- or 7-membered azacyclic group optionally containing one additional O, N or S atom in the azacycle and having 3-8 carbon atoms (including the carbon atoms contained in any optional substituent(s) of the azacycle));

- A and B are independently N or CH;
- $R_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$  and  $R_9$  are independently hydrogen or  $C_{1-4}$ alkyl;
- $R_{10}$  is a group of the formula (a) or (b):



wherein:

- Z is  $C_{1-4}$ alkyl, halo $C_{1-4}$ alkyl,  $C_{3-6}$ cycloalkyl, phenyl, heterocyclyl, a 5- or 6-membered heteroaromatic ring or a 8- to 11-membered bicyclic group, any of which is optionally substituted by 1, 2, 3 or 4 substituents selected from the group consisting of: halogen, hydroxy, oxo, cyano, nitro,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy, halo $C_{1-4}$ alkyl, halo $C_{1-4}$ alkoxy,  $C_{1-4}$ alkylenedioxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkylsulfonyl,  $C_{1-4}$ alkylsulfonyloxy, halo $C_{1-4}$ alkylsulfonyl, halo $C_{1-4}$ alkylsulfonyloxy,  $C_{1-4}$ alkylsulfinyl,  $C_{1-4}$ alkylthio,  $R_{17}SO_2N(R_{18})$ -,  $R_{17}R_{18}NSO_2$ -,  $R_{17}R_{18}N$ -,  $R_{17}R_{18}NCO$ -,  $R_{17}CONR_{18}$ - and a 5- or 6-membered heteroaromatic ring which is optionally substituted by one or two  $C_{1-2}$ alkyl, halo $C_{1-2}$ alkyl or  $R_{17}R_{18}N$ - (wherein  $R_{17}$  and  $R_{18}$  are independently hydrogen or  $C_{1-4}$ alkyl, or  $R_{17}$  and  $R_{18}$  together form  $C_{3-6}$ alkylene); and wherein substituents positioned *ortho* to one another may be linked to form a 5- or 6-membered ring; and
- $R_{15}$  and  $R_{16}$  are independently hydrogen or  $C_{1-4}$ alkyl and t is 1, 2, 3 or 4, or  $-(CR_{15}R_{16})_t$  forms a  $C_{3-6}$ cycloalkylene linker.

In formula (I), "-S-" means thio (sulfur).

The term " $C_{1-4}$ alkyl" refers to an alkyl group having from one to four carbon atoms, in all isomeric forms, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl and tert-butyl.

The term "alkylene" refers to a straight or branched chain divalent hydrocarbon radical. Examples of C<sub>1-3</sub>alkylene groups include methylene, ethylene and n-propylene. Examples of "C<sub>1-4</sub>alkylene" include, in addition to the above, n-butylene.

5

The term "C<sub>1-4</sub>alkoxy" refers to a straight chain or branched chain alkoxy (or "alkyloxy") group having from one to four carbon atoms, such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy and tert-butoxy.

- 10 The term "halogen" and its abbreviation "halo" refer to fluorine (F), chlorine (Cl), bromine (Br) or iodine (I). Where the term "halo" is used before another group, it indicates that the group is substituted by one, two or three halogen atoms. For example, "haloC<sub>1-4</sub>alkyl" refers to groups such as trifluoromethyl, bromoethyl, trifluoropropyl, and other groups derived from C<sub>1-4</sub>alkyl groups as defined above; and the term "haloC<sub>1-4</sub>alkoxy" refers to
- 15 groups such as trifluoromethoxy, bromoethoxy, trifluoropropoxy, and other groups derived from C<sub>1-4</sub>alkoxy groups as defined above.

- The term "C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl" refers to a C<sub>1-4</sub>alkoxy group attached through a C<sub>1-4</sub>alkylene group, for example methoxymethyl, ethoxymethyl, propoxyethyl, isopropoxyethyl and others derived from the C<sub>1-4</sub>alkoxy and C<sub>1-4</sub>alkyl groups as defined above.
- 20

The term "C<sub>1-4</sub>alkylthio" refers to a C<sub>1-4</sub>alkyl group attached through a sulfur atom (-S-). Examples of C<sub>1-4</sub>alkylthio include methylthio, ethylthio, propylthio and butylthio.

- 25 The term "C<sub>3-6</sub>cycloalkyl" refers to a cycloalkyl group having from three to six carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. The term "C<sub>3-6</sub>cycloalkylene" refers to a divalent cycloalkyl group, such as cyclopropylene, cyclobutylene, cyclopentylene and cyclohexylene.
- 30 The term "C<sub>3-6</sub>cycloalkylC<sub>1-4</sub>alkyl" refers to a cycloalkyl group attached through a C<sub>1-4</sub>alkylene group, such as cyclopropylmethyl, cyclobutylethyl, and others derived from C<sub>3-6</sub>cycloalkyl groups and C<sub>1-4</sub>alkyl groups as defined above.
- 35 The term "aryl" refers to phenyl or a 5- or 6-membered heteroaromatic ring. Examples of 5- or 6-membered heteroaromatic rings include furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, pyridinyl, triazolyl, triazinyl, pyridazyl, pyrimidinyl, isothiazolyl, isoxazolyl, pyrazinyl, pyrazolyl and pyrimidinyl.
- 40 The term "arylC<sub>1-4</sub>alkyl" refers to an aryl group attached through a C<sub>1-4</sub>alkylene group. The C<sub>1-6</sub>alkylene group may be in any suitable isomeric form. Examples of arylC<sub>1-4</sub>alkyl include benzyl, phenethyl (including phenyl-CH<sub>2</sub>CH<sub>2</sub>- and phenyl-C(CH<sub>3</sub>)-) and others derived from the aryl groups and C<sub>1-4</sub>alkyl groups as defined above.



The terms "arylC<sub>1-4</sub>alkoxy" refers to an aryl group attached through a C<sub>1-4</sub>alkoxy group. Examples of arylC<sub>1-4</sub>alkoxy include benzyloxy (phenyl-CH<sub>2</sub>O-) and phenylethoxy.

- 5 The term "sulfonyl" refers to the group -SO<sub>2</sub>-. Thus, the term "C<sub>1-4</sub>alkylsulfonyl" includes methylsulfonyl, ethylsulfonyl, and others derived from the C<sub>1-4</sub>alkyl groups defined above. The term "haloC<sub>1-4</sub>alkylsulfonyl" refers to groups such as trifluoromethanesulfonyl and pentafluoroethylsulfonyl. The term "arylsulfonyl" includes phenylsulfonyl, pyridinylsulfonyl, and others derived from aryls as defined above.

10

The term "arylcarboxamido" refers to groups such as phenylcarboxamido and pyridinylcarboxamido, and others derived from the aryl groups as defined above.

15

The term "C<sub>1-4</sub>alkylenedioxy" refers to groups such as methylenedioxy, ethylenedioxy and others derived from C<sub>1-4</sub>alkyl as defined above.

20

The term "5- or 6-membered heteroaromatic ring" refers to a monocyclic 5- or 6-membered heterocyclic group containing 1, 2, 3 or 4 heteroatoms, for example from 1 to 3 heteroatoms, selected from O, N and S. When the group contains 2-4 heteroatoms, one may be selected from O, N and S and the remaining heteroatoms may be N. Examples of 5 and 6-membered heteroaromatic rings include pyrrolyl, pyrrolinyl, pyrazolinyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, oxadiazolyl, isothiazolyl, thiazolyl, furyl, thienyl, thiadiazolyl, pyridyl, triazolyl, thiazinyl, triazinyl, pyridazinyl, pyrimidinyl and pyrazinyl.

25

30

The term "8- to 11-membered bicyclic group" refers to a bicyclic ring system containing a total of 8, 9, 10 or 11 carbon atoms, wherein 1, 2, 3 or 4 or 5 of the carbon atoms are optionally replaced by a heteroatom independently selected from O, S and N. The term includes bicyclic systems wherein both rings are aromatic, as well as bicyclic ring systems wherein one of the rings is partially or fully saturated. Examples of 8- to 11- membered bicyclic groups wherein both rings are aromatic include indenyl, naphthyl and azulenyl. Examples of 8- to 11-membered bicyclic groups having 1, 2, 3, 4 or 5 heteroatoms, in which both rings are aromatic, include: 6*H*-thieno[2,3-*b*]pyrrolyl, imidazo[2,1-*b*][1,3]thiazolyl, imidazo[5,1-*b*][1,3]thiazolyl, [1,3]thiazolo[3,2-*b*][1,2,4]triazolyl, indolyl, isoindolyl, indazolyl, benzimidazolyl e.g. benzimidazol-2-yl, benzoxazolyl e.g. benzoxazol-2-yl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzothienyl, benzofuranyl, naphthridinyl, quinolyl, quinoxalinyl, quinazolinyl, cinnolinyl and isoquinolyl. Examples of 8- to 11-membered bicyclic groups having 1, 2, 3, 4 or 5 heteroatoms, in which one of the rings is partially or fully saturated includes dihydrobenzofuranyl, indanyl, tetrahydronaphthyl, indolinyl, isoindolinyl, tetrahydroisoquinolyl, tetrahydroquinolyl, benzoxazinyl and benzoazepinyl.

40

The term "heterocyclyl" refers to a 5 or 6-membered monocyclic or 8 to 11-membered bicyclic group wherein 1, 2, 3, 4 or 5 of the carbon atoms are replaced by a heteroatom independently selected from O, S and N and which is partially or fully saturated. Examples of "heterocyclyl" which are fully saturated 5 or 6-membered monocyclic rings include pyrrolidinyl, imidazolidinyl, pyrazolidinyl, isothiazolyl, thiazolyl, tetrahydrofuranyl, dioxolanyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, tetrahydrothienyl, dioxanyl, tetrahydro-2*H*-pyranyl and dithianyl. Examples of "heterocyclyl" groups which are partially saturated 5 or 6-membered monocyclic rings include oxazoliny, isoaxazoliny, imidazoliny, pyrazoliny, 1,2,3,6-tetrahydropyridyl and 3,6-dihydro-2*H*-pyranyl. Examples of "heterocyclyl" groups which are fully saturated 8 to 11-membered bicyclic rings include decahydroquinoliny, octahydro-2*H*-1,4-benzoxazinyl and octahydro-1*H*-cyclopenta[*b*]pyridiny. Examples of "heterocyclyl" groups which are partially saturated 8 to 11-membered bicyclic rings include 2,3-dihydro-1*H*-indolyl, 1,2,3,4-tetrahydroquinoliny, 1,2,3,4-tetrahydroisoquinoliny and 2,3,4,5-tetrahydro-1*H*-3-benzazepiny.

Any of these groups may be attached to the rest of the molecule at any suitable position.

As used herein, the term "salt" refers to any salt of a compound according to the present invention prepared from an inorganic or organic acid or base, quaternary ammonium salts and internally formed salts. Physiologically acceptable salts are particularly suitable for medical applications because of their greater aqueous solubility relative to the parent compounds. Such salts must clearly have a physiologically acceptable anion or cation. Suitably physiologically acceptable salts of the compounds of the present invention include acid addition salts formed with inorganic acids such as hydrochloric, hydrobromic, hydroiodic, phosphoric, metaphosphoric, nitric and sulfuric acids, and with organic acids, such as tartaric, acetic, trifluoroacetic, citric, malic, lactic, fumaric, benzoic, formic, propionic, glycolic, gluconic, maleic, succinic, camphorsulfuric, isothionic, mucic, gentisic, isonicotinic, saccharic, glucuronic, furoic, glutamic, ascorbic, anthranilic, salicylic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, pantothenic, stearic, sulfinilic, alginic, galacturonic and arylsulfonic, for example benzenesulfonic and *p*-toluenesulfonic, acids; base addition salts formed with alkali metals and alkaline earth metals and organic bases such as *N,N*-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (*N*-methylglucamine), lysine and procaine; and internally formed salts. Salts having a non-physiologically acceptable anion or cation are within the scope of the invention as useful intermediates for the preparation of physiologically acceptable salts and/or for use in non-therapeutic, for example, *in vitro*, situations.

When  $R_2$  or  $R_3$  contains an aryl moiety, *i.e.*  $R_2$  or  $R_3$  is aryl, arylC<sub>1-4</sub>alkoxy, aryloxy, arylthio, arylmethyl, aroyl, aryloxymethyl, arylsulfonyl, aryl-NR'-, arylsulfonyloxy, arylsulfonylC<sub>1-4</sub>alkyl, arylsulfonamido, arylcarboxamido, arylsulfonamidoC<sub>1-4</sub>alkyl, arylcarboxamidoC<sub>1-4</sub>alkyl, aroylC<sub>1-4</sub>alkyl or arylC<sub>1-4</sub>alkanoyl, the aryl moiety is optionally substituted by one or two substituents selected from: halogen, cyano, C<sub>1-2</sub>alkyl (e.g. methyl), fluoroC<sub>1-2</sub>alkyl (eg

trifluoromethyl), C<sub>1-2</sub>alkoxy (e.g. methoxy), C<sub>1-2</sub>alkylenedioxy (e.g. methylenedioxy), C<sub>1-3</sub>alkanoyl (e.g. acetyl), C<sub>2</sub>alkanoylamino (e.g. acetylamino), fluoroC<sub>1</sub>alkylsulfonyl (e.g. trifluoromethylsulfonyl) and methylsulfonyl. For example, the aryl moiety is optionally substituted by methyl.

5

When R<sub>2</sub> or R<sub>3</sub> is a group R<sub>11</sub>CON(R<sub>12</sub>)(CH<sub>2</sub>)<sub>r</sub>, R<sub>11</sub>R<sub>12</sub>NCO(CH<sub>2</sub>)<sub>r</sub> or R<sub>11</sub>R<sub>12</sub>NSO<sub>2</sub>(CH<sub>2</sub>)<sub>r</sub> and R<sub>11</sub>CONR<sub>12</sub> or R<sub>11</sub>R<sub>12</sub>N together form a 4-, 5-, 6- or 7-membered azacyclic group, then this is characterised by: (i) containing one additional O, N or S atom in the azacycle, for example the azacyclic group being 1,4-morpholin-4-yl and/or (ii) having 1 or 2 optional C<sub>1-2</sub>alkyl substituents whose carbon atoms are included in the azacyclic group's 3-8 carbon atoms. One, two or more F atoms can optionally be included as substituents of the carbon atoms of the heterocycle. The term "azacyclic group" should be interpreted to cover only stable azacycles such as 1,4-morpholine and piperazine and not for example 1,3-morpholine. Saturated azacycles, in particular piperidinyl, pyrrolidinyl, 1,4-morpholinyl, and including the corresponding α-oxo-azacycles R<sub>11</sub>CONR<sub>12</sub>, are preferred.

10

15

In one embodiment, R<sub>2</sub> or R<sub>3</sub> is halogen, cyano, acetyl, trifluoromethyl, pentafluoroethyl, trifluoromethoxy, C<sub>1-4</sub>alkylsulfonyl, C<sub>1-4</sub>alkylsulfonyloxy, R<sub>11</sub>R<sub>12</sub>NSO<sub>2</sub> (where each of R<sub>11</sub> and R<sub>12</sub> is independently hydrogen or C<sub>1-4</sub>alkyl or R<sub>11</sub>R<sub>12</sub>N together form a 4-, 5-, 6- or 7-membered azacyclic group optionally containing one additional O, N or S atom in the azacycle and having 3-8 carbon atoms), a heterocyclyl, or a 5- or 6-membered heteroaromatic ring which is optionally substituted by one or two substituents selected from: halogen, cyano, C<sub>1-2</sub>alkyl (e.g. methyl), haloC<sub>1-2</sub>alkyl (e.g. trifluoromethyl), C<sub>1-2</sub>alkoxy (e.g. methoxy), C<sub>1-2</sub>alkylenedioxy (e.g. methylenedioxy), C<sub>1-3</sub>alkanoyl (e.g. acetyl), C<sub>2</sub>alkanoylamino (e.g. acetylamino), haloC<sub>1</sub>alkylsulfonyl (e.g. trifluoromethylsulfonyl) and methylsulfonyl.

20

25

In one embodiment, R<sub>3</sub> is hydrogen.

30

Examples of R<sub>2</sub> include: C<sub>1-4</sub>alkyl, haloC<sub>1-4</sub>alkyl, halogen, C<sub>1-4</sub>alkylsulfonyl (e.g. methylsulfonyl or ethylsulfonyl), haloC<sub>1-4</sub>alkylsulfonyl (e.g. trifluoromethylsulfonyl), C<sub>1-4</sub>alkylsulfonyloxy (e.g. methylsulfonyloxy), haloC<sub>1-4</sub>alkylsulfonyloxy (e.g. trifluoromethylsulfonyloxy), R<sub>11</sub>R<sub>12</sub>NSO<sub>2</sub> (where each of R<sub>11</sub> and R<sub>12</sub> is independently hydrogen or C<sub>1-4</sub>alkyl or R<sub>11</sub>R<sub>12</sub>N together form a 4-, 5-, 6- or 7-membered azacyclic group optionally containing one additional O, N or S atom in the azacycle and having 3-8 carbon atoms, e.g. a piperidin-1-ylsulfonyl, pyrrolidin-1-ylsulfonyl or 1,4-morpholin-4-ylsulfonyl), a 5- or 6-membered heteroaromatic or a heterocyclyl, each of which is optionally substituted by one or two substituents selected from: halogen, cyano, C<sub>1-2</sub>alkyl (e.g. methyl or trifluoromethyl), C<sub>1-2</sub>alkoxy (e.g. methoxy), C<sub>1-2</sub>alkylenedioxy (e.g. methylenedioxy), C<sub>1-3</sub>alkanoyl (e.g. acetyl), C<sub>2</sub>alkanoylamino (e.g. acetylamino), haloC<sub>1</sub>alkylsulfonyl (e.g. trifluoromethylsulfonyl) and methylsulfonyl.

35

40

Suitably,  $R_2$  is bromo, cyano, hydroxy, chloro, methoxy, tert-butyl, methylsulfonyl, ethylsulfonyl, N,N-dimethylaminosulfonyl, pyrrolidin-1-ylsulfonyl, 1,4-morpholin-4-ylsulfonyl, methylsulfonyloxy, pyrazin-2-yl, 5-methyl-oxazol-2-yl or 5-methyl-isoxazol-3-yl.

- 5 In one embodiment, at least one of  $R_1$  and  $R_4$  is hydrogen. For example, both  $R_1$  and  $R_4$  are hydrogen, or all of  $R_1$ ,  $R_3$  and  $R_4$  are hydrogen.

In one embodiment, at least one of A and B is nitrogen. For example, A and B may both be nitrogen.

10

In one embodiment,  $R_5$ ,  $R_6$ ,  $R_7$  and  $R_8$  are all hydrogen.

In one embodiment,  $R_9$  is methyl.

- 15  $R_{10}$  may be formula (a) or (b). For formula (a) and (b), in one embodiment, Z may be optionally substituted phenyl, an optionally substituted bicyclic group such as quinolinyl (e.g. 2-, 3-, 4-, 5- or 6-quinolinyl), 4-tetrahydro-2H-pyranyl, furyl (e.g. 2-furyl), thienyl (e.g. 2-thienyl), pyridyl (e.g. 4-pyridyl), indolyl, pyrazolopyrimidyl (e.g. pyrazolo[1,5-a]pyrimidyl), cinnolinyl, benzo[b]furanyl, thienopyridine or pyrrolopyridyl. Examples of Z include 4-  
20 tetrahydro-2H-pyranyl, 4-trifluoromethylphenyl, furyl e.g. 2-furyl, thienyl e.g. 2-thienyl, pyridyl e.g. 4-pyridyl, 2-methylquinolinyl and 4-methyl,3-oxazol-5-yl.


- When  $R_{10}$  is a group of formula (b), and  $R_{12}$  and  $R_{13}$  are independently hydrogen or  $C_{1-4}$ alkyl and t is 1, 2, 3 or 4, examples include  $-(CH_2)_t-Z$ , and  $-(CHCH_3)_t-Z$ . When the group -  
25  $(CR_{15}R_{16})_t$  in formula (b) forms a  $C_{3-6}$ cycloalkylene linker, examples include groups such as:



- 30 In one embodiment, Z is unsubstituted or substituted by one or more substituents selected from: halogen, or cyano,  $C_{1-2}$ alkyl (e.g. methyl), halo $C_{1-2}$ alkyl (e.g. trifluoromethyl),  $C_{1-2}$ alkoxy (e.g. methoxy), halo $C_{1-4}$ alkoxy (e.g. trifluoromethoxy),  $C_{1-2}$ alkylenedioxy (e.g. methylenedioxy),  $C_{2-3}$ alkanoyl (e.g. acetyl),  $C_2$ alkanoylamino (e.g. acetylamino), methylsulfonyl, halo $C_1$ alkylsulfonyl (e.g. trifluoromethylsulfonyl),  $C_1$ alkylsulfonyloxy (e.g. methylsulfonyloxy),  $C_1$ alkylaminosulfonyl (e.g. methylaminosulfonyl),  $C_1$ alkylsulfonylamino (e.g. methylsulfonylamino) and  $C_1$ alkylaminocarbonyl (e.g. methylaminocarbonyl).  
35

- In one embodiment,  $R_{10}$  is a group of formula (a) as defined in formula (I). For example,  $R_{10}$  may be optionally substituted phenyl, such as unsubstituted phenyl or fluorophenyl  
40 (e.g. 4-fluorophenyl), or optionally substituted quinolinyl (e.g. 6-quinolinyl).

In one embodiment, a compound of formula (IA) or a salt thereof is provided:



(IA)

wherein A, B and R<sub>9</sub> are as defined for formula (I);

X is a 5- or 6-membered heteroaromatic ring optionally substituted by 1, 2 or 3 substituents selected from the group consisting of: halogen, cyano, C<sub>1-2</sub>alkyl, fluoroC<sub>1-2</sub>alkyl, C<sub>1-2</sub>alkoxy, C<sub>1-3</sub>alkanoyl, C<sub>2</sub>alkanoylamino, fluoroC<sub>1</sub>alkylsulfonyl and methylsulfonyl; and

Y is heterocyclyl, a 5- or 6-membered heteroaromatic ring or a 8- to 11-membered bicyclic group, any of which is optionally substituted by 1, 2, 3 or 4 substituents selected from the group consisting of: halogen, cyano, C<sub>1-2</sub>alkyl, haloC<sub>1-2</sub>alkyl, C<sub>1-2</sub>alkoxy, haloC<sub>1-2</sub>alkoxy, C<sub>1-2</sub>alkylenedioxy, C<sub>2-3</sub>alkanoyl, C<sub>2</sub>alkanoylamino, methylsulfonyl, haloC<sub>1</sub>alkylsulfonyl, methylsulfonyloxy, methylaminosulfonyl, methylsulfonylamino and methylaminocarbonyl.

All embodiments and features of formula (I) apply to formula (IA).

Example compounds of the present invention include:

1. 7-(5-Methyl-3-isoxazolyl)-3-(2-([4-methyl-1,3-oxazol-5yl)-4H-1,2,4-triazol-3-yl]thio)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride
2. 7-(5-Methyl-3-isoxazolyl)-3-(2-([4-methyl-5-(tetrahydro-2H-pyran-4-yl)-4H-1,2,4-triazol-3-yl]thio)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride
3. 7-(5-Methyl-3-isoxazolyl)-3-(2-([4-methyl-5-(2-methyl-5-quinolinyl)-4H-1,2,4-triazol-3-yl]thio)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride
4. 7-(5-Methyl-3-isoxazolyl)-3-(2-([4-methyl-5-(2-methyl-6-quinolinyl)-4H-1,2,4-triazol-3-yl]thio)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride

and salts thereof.

It will be appreciated that for use in medicine the salts of the compounds of the invention should be pharmaceutically (i.e. physiologically) acceptable. Suitable pharmaceutically acceptable salts will be apparent to those skilled in the art and include for example acid addition salts formed with inorganic acids e.g. hydrochloric, hydrobromic, sulfuric, nitric or phosphoric acid; and organic acids e.g. succinic, maleic, acetic, fumaric, citric, tartaric, benzoic, p-toluenesulfonic, methanesulfonic or naphthalenesulfonic acid. Other non-pharmaceutically acceptable salts eg. oxalates, may be used, for example in the isolation

of compounds of the invention and are included within the scope of this invention. Also included within the scope of the invention are solvates, hydrates, complexes and prodrugs of compounds of the invention.

- 5 Certain of the compounds of the invention may form acid addition salts with one or more equivalents of the acid. The present invention includes within its scope all possible stoichiometric and non-stoichiometric forms.

- 10 Certain groups/substituents included in the present invention may be present as isomers. The present invention includes within its scope all such isomers, including racemates, enantiomers, tautomers and mixtures thereof. Certain of the substituted heteroaromatic rings included in compounds of formula (I) may exist in one or more tautomeric forms. The present invention includes within its scope all such tautomeric forms, including mixtures.


- 15 Preferred compounds have a molecular weight of 800 or less. Still more preferred are compounds having a molecular weight of 600 or less. Generally, and without being limited thereto, such compounds may have higher oral bioavailability, and sometimes higher solubility and/or brain penetrancy. Molecular weight here refers to that of the unsolvated  
20 free base compound, excluding any molecular weight contributed by addition salts, solvent (e.g. water) molecules, prodrug molecular parts cleaved off *in vivo*, etc.

- In general, the compounds or salts of the invention should be interpreted as excluding those compounds (if any) which are so chemically unstable, either per se or in water, that  
25 they are clearly unsuitable for pharmaceutical use through all administration routes, whether oral, parenteral or otherwise. Such compounds are known to the skilled chemist. Prodrugs or compounds which are stable *ex vivo* and which are convertible in the mammalian (e.g. human) body to the inventive compounds are however included.

- 30 The present invention also provides a process for preparing a compound of formula (I), which process comprises:


(a) reacting a compound of formula (II):

35



(II)

wherein  $R_1$  to  $R_8$  are as defined for formula (I) and L is a leaving group; with a compound of formula (III):




(III)

wherein A, B,  $R_9$  and  $R_{10}$  are as defined for formula (I); or

5


(b) for a compound of formula (I) wherein  $R_2$  is aryl, reacting a compound of formula (IV):



(IV)

10 wherein  $R_1$ ,  $R_3$  to  $R_{10}$ , A and B are as defined for formula (I) and W is halogen or a trifluoromethylsulfonyloxy group, or W is a group M selected from a boron derivative (e.g. a boronic acid function  $B(OH)_2$ ) or a metal function such as trialkylstannyl (e.g.  $SnBu_3$ ), zinc halide or magnesium halide; with a compound aryl- $W^1$ , wherein aryl is as defined for formula (I),  $W^1$  is halogen or a trifluoromethylsulfonyloxy group when W is a group M or  $W^1$  is a group M as defined above when W is halogen or a trifluoromethylsulfonyloxy group; or

15 (c) for a compound of formula (I) wherein  $R_2$  is aryloxy or arylthio, reacting a compound of formula (V):



wherein G is oxygen or sulfur, and  $R_1$ ,  $R_3$  to  $R_{10}$ , A and B are as defined for formula (I); with a reagent serving to introduce the aryl group;

5

and optionally thereafter for any of the steps (a), (b) or (c):

- removing any protecting group(s); and/or
- forming a salt; and/or
- converting one compound of formula (I) to a different compound of formula (I).

10

Process (a) may be effected using conventional methods for the formation of a thioether. The leaving group L can be halogen such as chlorine. Alternatively L can be a sulfonyloxy group such  $C_{1-4}$ alkylsulfonyloxy (e.g. methanesulfonyloxy or trifluoromethanesulfonyloxy); or  $AR'$ -sulfonyloxy wherein  $AR'$  is optionally substituted phenyl, an optionally substituted 5- or 6-membered aromatic heterocyclic group, or an optionally substituted bicyclic group, preferably optionally substituted phenyl, wherein in each case the optional substituents are one or more  $C_{1-2}$ alkyl groups; e.g. *para*-toluenesulfonyloxy. When L is a halogen the reaction may be carried out using a base such as lithium hydroxide in a solvent such as *N,N*-dimethylformamide.

20

The reaction in process (b), and the reaction in process (d), may be effected in the presence of a transition metal e.g., palladium catalyst such as *bis*-triphenylphosphinepalladium dichloride or *tetrakis*-triphenylphosphinepalladium (0). When M is a boronic acid function such as  $B(OH)_2$  the reaction may be carried out under basic conditions, for example using aqueous sodium carbonate in a suitable solvent such as dioxane. When M is trialkylstannyl the reaction may be carried out in an inert solvent, such as xylene or dioxane optionally in the presence of LiCl. When M is a zinc or magnesium halide the reaction may be effected in an aprotic solvent such as tetrahydrofuran. The substituent W is preferably halogen such as bromine, or a sulfonyloxy group such as trifluoromethylsulfonyloxy; and  $W^1$  is preferably a group M, such as trialkylstannyl or  $B(OH)_2$ .

25


30

35

In process (c), the reagent serving to introduce the aryl group is preferably a compound of formula aryl-Hal, wherein Hal is halogen. The reaction may be effected in the presence of a base, such as potassium carbonate, in a solvent such as *N,N*-dimethylformamide.



A compound of formula (II) may itself be prepared by reacting a compound of formula (VII):



**Formula (VII)**

wherein  $R^1$  to  $R^4$  are as hereinbefore defined; with a compound of formula (VIII):



**Formula (VIII)**

wherein L is as herein defined and L' is a leaving group, e.g., a bromine atom or alternatively with a compound of formula (IX)



**Formula (IX)**

wherein L is as herein defined, in the presence of a hydride source such as sodium triacetoxyborohydride.

Compounds of formula (I) may be converted to another compound of formula (I) by suitable methods known to the skilled person, such as:

- (i) converting one or more of  $R_1$  to  $R_4$  from alkoxy (e.g. methoxy) to hydroxyl; and
- (ii) converting one or more of  $R_2$  or  $R_3$  from hydroxy to sulfonyloxy, such as alkylsulfonyloxy e.g. methanesulfonyloxy or trifluoromethanesulfonyloxy.

Compounds of formula (I) have been found to exhibit affinity for dopamine receptors, in particular the  $D_3$  receptor, and are expected to be useful in the treatment of disease states which require modulation of such receptors, such as psychotic conditions. Many of the compounds of formula (I) have also been found to have greater affinity for dopamine  $D_3$  than for  $D_2$  receptors. Compounds for formula (I) have also been found to exhibit low affinity for the  $H1$  receptor. A low affinity for the  $H1$  receptor generally leads to avoidance of: (1) sedation, somnolence, and fatigue; (2) cardiotoxicity; (3) potentiation of opioid-induced sedation and respiratory depression; (4) short-term weight gain; (5) impaired cognition (memory, spatial cognition, attention, tracking performance); (6) impaired psychomotor performance including quick tolerance to these effects, and (7) altered neuroendocrine regulation of prolactin and potentially other hormones.

The therapeutic effect of currently available antipsychotic agents (neuroleptics) is generally believed to be exerted via blockade of D<sub>2</sub> receptors; however this mechanism is also thought to be responsible for undesirable extrapyramidal side effects (eps) associated with many neuroleptic agents. Without wishing to be bound by theory, it has been suggested that blockade of the recently characterised dopamine D<sub>3</sub> receptor may give rise to beneficial antipsychotic activity without significant eps. (see for example Sokoloff et al, Nature, 1990; 347: 146-151; and Schwartz et al, Clinical Neuropharmacology, Vol 16, No. 4, 295-314, 1993). Preferred compounds of the present invention are therefore those which have higher (e.g.  $\geq 10\times$  or  $\geq 100\times$  higher) affinity for dopamine D<sub>3</sub> than dopamine D<sub>2</sub> receptors (such affinity can be measured using standard methodology for example using cloned dopamine receptors – see herein). Said compounds may advantageously be used as selective modulators of D<sub>3</sub> receptors.

Compounds of formula (I) will be used for treatment of all aspects of drug dependency including withdrawal symptoms from drugs of abuse such as alcohol, cocaine, opiates, nicotine, benzodiazepines and inhibition of tolerance induced by opioids. In addition, compounds of formula (I) and pharmaceutically acceptable salts and solvates thereof will be used to reduce craving and therefore will be useful in the treatment of drug craving. Drug craving can be defined as the incentive motivation to self-administer a psychoactive substance that was previously consumed. Three main factors are involved in the development and maintenance of drug craving: (1) Dysphoric states during drug withdrawal can function as a negative reinforcer leading to craving; (2) Environmental stimuli associated with drug effects can become progressively more powerful (sensitization) in controlling drug seeking or craving, and (3) A cognition (memory) of the ability of drugs to promote pleasurable effects and to alleviate a dysphoric state during withdrawal. Craving may account for the difficulty that individuals have in giving up drugs of abuse and therefore contributes significantly to the development and maintenance of drug dependence.

The compounds of formula (I) are of potential use as antipsychotic agents for example in the treatment of schizophrenia, schizo-affective disorders, psychotic depression, mania, paranoid and delusional disorders. Furthermore, they could have utility as adjunct therapy in Parkinsons Disease, particularly with compounds such as L-DOPA and possibly dopaminergic agonists, to reduce the side effects experienced with these treatments on long term use (e.g. see Schwartz et al., Brain Res. Reviews, 1998, 26, 236-242). From the localisation of D<sub>3</sub> receptors, it could also be envisaged that the compounds could also have utility for the treatment of substance abuse where it has been suggested that D<sub>3</sub> receptors are involved (e.g. see Levant, 1997, Pharmacol. Rev., 49, 231-252). Examples of such substance abuse include alcohol, cocaine, heroin and nicotine abuse. Other conditions which may be treated by the compounds include dyskinetic disorders such as Parkinson's disease, neuroleptic-induced parkinsonism and tardive dyskinesias; depression; anxiety, cognitive impairment including memory disorders such as Alzheimers disease, eating disorders, sexual dysfunction, sleep disorders, emesis, movement

disorders, obsessive-compulsive disorders, amnesia, aggression, autism, vertigo, dementia, circadian rhythm disorders and gastric motility disorders e.g. IBS.

5 In a further aspect therefore the present invention provides a method of treating a condition for which modulation (especially antagonism/inhibition) of dopamine receptors (especially dopamine D<sub>3</sub> receptors) is beneficial, which comprises administering to a mammal (e.g. human) in need thereof an effective amount of a compound of formula (I) or a pharmaceutically (i.e physiologically) acceptable salt thereof. Such conditions in particular include psychoses/psychotic conditions such as schizophrenia, and substance  
10 abuse and/or drug dependency.

The invention also provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for the treatment of a condition in a mammal for which modulation (especially antagonism/inhibition) of  
15 dopamine receptors (especially dopamine D<sub>3</sub> receptors) is beneficial.

The invention also provides a compound of formula (I) or a pharmaceutically acceptable salt thereof for use in the treatment of a condition in a mammal for which modulation (especially antagonism/inhibition) of dopamine receptors (especially dopamine D<sub>3</sub>  
20 receptors) is beneficial.

In one embodiment, D<sub>3</sub> antagonists according to the present invention are used in the treatment of psychoses such as schizophrenia or in the treatment of substance abuse and/or drug dependency.  
25

Thus, a still further aspect the invention provides a method of treating a psychotic condition (e.g. schizophrenia) or substance abuse and/or drug dependency which comprises administering to a mammal (e.g. human) in need thereof an effective amount of a compound of formula (I) as herein defined or a pharmaceutically acceptable salt thereof.  
30

Also provided is the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for the treatment of a psychotic condition (e.g. schizophrenia) or substance abuse and/or drug dependency in a mammal.  
35

Also provided is a compound of formula (I) or a pharmaceutically acceptable salt thereof for use in the treatment of a psychotic condition (e.g. schizophrenia) or substance abuse and/or drug dependency in a mammal.

40 Also provided is a compound of formula (I) or a pharmaceutically acceptable salt thereof for use as an active therapeutic substance in a mammal, e.g. for use in the treatment of any of the conditions described herein.

"Treatment" includes prophylaxis, where this is appropriate for the relevant condition(s).

For use in medicine, the compounds of the present invention are usually administered as a standard pharmaceutical composition. The present invention therefore provides in a further aspect a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically (i.e physiologically) acceptable salt thereof and a pharmaceutically (i.e physiologically) acceptable carrier. The pharmaceutical composition can be for use in the treatment of any of the conditions described herein.

10 The compounds of formula (I) may be administered by any convenient method, for example by oral, parenteral (e.g. intravenous), buccal, sublingual, nasal, rectal or transdermal administration and the pharmaceutical compositions adapted accordingly.

15 The compounds of formula (I) and their pharmaceutically acceptable salts which are active when given orally can be formulated as liquids or solids, for example syrups, suspensions or emulsions, tablets, capsules and lozenges.

20 A liquid formulation will generally consist of a suspension or solution of the compound or pharmaceutically acceptable salt in a suitable liquid carrier(s) for example an aqueous solvent such as water, ethanol or glycerine, or a non-aqueous solvent, such as polyethylene glycol or an oil. The formulation may also contain a suspending agent, preservative, flavouring or colouring agent.

25 A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and cellulose.

30 A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule.

35 Typical parenteral compositions consist of a solution or suspension of the compound or pharmaceutically acceptable salt in a sterile aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration.

40 Compositions for nasal administration may conveniently be formulated as aerosols, drops, gels and powders. Aerosol formulations typically comprise a solution or fine suspension of the active substance in a pharmaceutically acceptable aqueous or non-aqueous

solvent and are usually presented in single or multidose quantities in sterile form in a sealed container, which can take the form of a cartridge or refill for use with an atomising device. Alternatively the sealed container may be a unitary dispensing device such as a single dose nasal inhaler or an aerosol dispenser fitted with a metering valve which is intended for disposal once the contents of the container have been exhausted. Where the dosage form comprises an aerosol dispenser, it will contain a propellant which can be a compressed gas such as compressed air or an organic propellant such as a fluoro-chlorohydrocarbon. The aerosol dosage forms can also take the form of a pump-atomiser.

Compositions suitable for buccal or sublingual administration include tablets, lozenges and pastilles, wherein the active ingredient is formulated with a carrier such as sugar and acacia, tragacanth, or gelatin and glycerin.

Compositions for rectal administration are conveniently in the form of suppositories containing a conventional suppository base such as cocoa butter.

Compositions suitable for transdermal administration include ointments, gels and patches.

In one embodiment, the composition is in unit dose form such as a tablet, capsule or ampoule.

Each dosage unit for oral administration contains for example from 1 to 250 mg (and for parenteral administration contains for example from 0.1 to 25 mg) of a compound of the formula (I) or a pharmaceutically acceptable salt thereof calculated as the free base.

The pharmaceutically acceptable compounds of the invention will normally be administered in a daily dosage regimen (for an adult patient) of, for example, an oral dose of between 1 mg and 500 mg, for example between 10 mg and 400 mg, e.g. between 10 and 250 mg or an intravenous, subcutaneous, or intramuscular dose of between 0.1 mg and 100 mg, for example between 0.1 mg and 50 mg, e.g. between 1 and 25 mg of the compound of the formula (I) or a pharmaceutically acceptable salt thereof calculated as the free base, the compound being administered 1 to 4 times per day. Suitably the compounds will be administered for a period of continuous therapy, for example for a week or more.

## Biological Test Methods

### Binding experiments on cloned dopamine (e.g. D2, D3 and D4) receptors

The ability of the compounds to bind selectively to human D2/D3/D4 dopamine receptors can be demonstrated by measuring their binding to cloned receptors. The inhibition constants ( $K_i$ ) of test compounds for displacement of [ $^{125}$ I]-iodosulpride binding to human D2/D3 and [ $^3$ H]-YM-09151 to D4 dopamine receptors expressed in CHO cells

were determined as follows. The cell lines were shown to be free from bacterial, fungal and mycoplasmal contaminants, and stocks of each were stored frozen in liquid nitrogen. Cultures were grown as monolayers or in suspension in standard cell culture media. Cells were recovered by scraping (from monolayers) or by centrifugation (from suspension cultures), and were washed two or three times by suspension in phosphate buffered saline followed by collection by centrifugation. Cell pellets were stored frozen at -80°C. Crude cell membranes were prepared by homogenisation followed by high-speed centrifugation, and characterisation of cloned receptors achieved by radioligand binding.

10 Preparation of CHO cell membranes: Cell pellets were gently thawed at room temperature, and resuspended in about 20 volumes of ice-cold Extraction buffer; 5mM EDTA, 50mM Trizma pre-set crystals (pH7.4@37°C), 1mM MgCl<sub>2</sub>, 5mM KCl and 120mM NaCl. The suspension was homogenised using an Ultra-Turrax at full speed for 15 seconds. The homogenate was centrifuged at 18,000 r.p.m for 15 min at 4°C in a Sorvall RC5C centrifuge. Supernatant was discarded, and homogenate re-suspended in extraction buffer then centrifugation was repeated. The final pellet was resuspended in 50mM Trizma pre-set crystals (pH 7.4 @ 37°C) and stored in 1ml aliquot tubes at -80°C (D2 = 3.0E+08 cells, D3 = 7.0E+07 cells and D4 = 1.0E+08 cells). The protein content was determined using a BCA protocol and bovine serum albumin as a standard (Smith, P. K., et al., Measurement of protein using bicinchoninic acid. Anal. Biochem. 150, 76-85 (1985)).

25 Binding experiments: Crude D2/D3 cell membranes were incubated with 0.03nM [<sup>125</sup>I]-Iodosulpride (~2000 Ci/mmol; Amersham, U. K.) and D4 with 0.8nM [<sup>3</sup>H]-YM-09151 (~85Ci/mmol; NEN, UK), and the test compound in a buffer containing 50mM Trizma pre-set crystals (pH 7.4 @ 37°C), 120mM NaCl, 5mM KCl, 2mM CaCl<sub>2</sub>, 1mM MgCl<sub>2</sub>, 0.3% (w/v) bovine serum albumin. The total volume is 0.2ml and incubated in a water bath at 37°C for 40 minutes. Following incubation, samples were filtered onto GF/B Unifilters using a Canberra Packard Filtermate, and washed four times with ice-cold 50mM Trizma pre-set crystals (pH 7.4 @ 37°C). The radioactivity on the filters was measured using a Canberra Packard Topcount Scintillation counter. Non-specific binding was defined with 10µM SKF-102161 (YM-09151). For competition curves, 10 serial log concentrations of competing cold drug were used (Dilution range: 10µM-10pM). Competition curves were analysed using Inflexion, an iterative curve fitting programme in Excel. Results were expressed as pKi values where  $pKi = -\log_{10}[Ki]$ .

The exemplified compounds have pKi values within the range of 7.5 - 9.5 at the dopamine D3 receptor. pKi results are only estimated to be accurate to about ±0.3-0.5.

#### 40 **Functional Activity at cloned dopamine receptors**

The functional activity of compounds at human D2 and human D3 receptors (i.e. agonism or antagonism) may be determined using a Cytosensor Microphysiometer (McConnell HM et al Science 1992 257 1906-1912). In Microphysiometer experiments, cells (hD2\_CHO

or hD3\_CHO) were seeded into 12mm Transwell inserts (Costar) at 300000 cells/cup in foetal calf serum (FCS)-containing medium. The cells were incubated for 6h at 37°C in 5%CO<sub>2</sub>, before changing to FCS-free medium. After a further 16-18h, cups were loaded into the sensor chambers of the Cytosensor Microphysiometer (Molecular Devices) and the chambers perfused with running medium (bicarbonate-free Dulbecco's modified Eagles medium containing 2 mM glutamine and 44 mM NaCl) at a flow rate of 100 ul/min. Each pump cycle lasted 90s. The pump was on for the first 60s and the acidification rate determined between 68 and 88s, using the Cytosoft programme. Test compounds were diluted in running medium. In experiments to determine agonist activity, cells were exposed (4.5 min for hD2, 7.5 min for hD3) to increasing concentrations of putative agonist at half hour intervals. Seven concentrations of the putative agonist were used. Peak acidification rate to each putative agonist concentration was determined and concentration-response curves fitted using Robofit [Tilford, N.S., Bowen, W.P. & Baxter, G.S. Br. J. Pharmacol. (1995), Vol. 115, 160P]. In experiments to determine antagonist potency, cells were treated at 30 min intervals with five pulses of a submaximal concentration of quinpirole (100 nM for hD2 cells, 30 nM for hD3 cells), before exposure to the lowest concentration of putative antagonist. At the end of the next 30 min interval, cells were pulsed again with quinpirole (in the continued presence of the antagonist) before exposure to the next highest antagonist concentration. In all, five concentrations of antagonist were used in each experiment. Peak acidification rate to each agonist concentration was determined and concentration-inhibition curves fitted using Robofit.


#### Human Histamine H1 receptor activity

Activity at the human Histamine H1 receptor can be measured using the general culture and assay conditions described in, for example, Smart et al, British Journal of Pharmacology (1999) 128, 1-3.

#### Examples

The invention is further illustrated by the following non-limiting examples.

#### Preparation 1: 3-(2-Chloroethyl)-7-(5-methyl-3-isoxazolyl)-2,3,4,5-tetrahydro-1H-3-benzazepine




To a solution of 7-(5-methyl-3-isoxazolyl)-2,3,4,5-tetrahydro-1H-3-benzazepine (WO0240471A2) (0.21 g, 0.92 mol) in a mixture of dichloroethane (2 mL) and acetonitrile (2 mL) at room temperature was added chloroacetaldehyde (0.23 mL) followed by sodium triacetoxyborohydride (0.39 g). The reaction mixture was stirred at room temperature for 1 h. Solvent was removed *in vacuo* and the residue was dissolved in water (5 mL) and CH<sub>2</sub>Cl<sub>2</sub> (5 mL). The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3X5 mL) and the organic layer was

dried over Na<sub>2</sub>SO<sub>4</sub>. Filtration and evaporation gave the crude product which was purified by flash chromatography (silica gel/ethylacetate:cyclohexanes 2:8) to give 0.11 g of the title compound (41% yield).

- 5 **NMR (<sup>1</sup>H, CDCl<sub>3</sub>):** δ 7.5 (m, 1H), 7.45 (m, 1H), 7.15 (m, 1H), 6.2 (s, 1H), 3.60-3.55 (m, 2H), 2.95-2.85 (m, 6H), 2.75-2.70 (m, 4H), 2.45 (s, 3H).

**Example 1: 7-(5-Methyl-3-isoxazolyl)-3-(2-([4-methyl-1,3-oxazol-5yl)-4H-1,2,4-triazol-3-yl]thio)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride**




10

- To a solution of 3-(2-chloroethyl)-7-(5-methyl-3-isoxazolyl)-2,3,4,5-tetrahydro-1H-3-benzazepine (0.03 g, 0.10 mmol) in dry DMF (0.5 mL), 4-methyl-5-(4-methyl-1,3-oxazol-5-yl)-4H-1,2,4-triazole-3-thiol (WO0240471A2) (0.10 mmol) was added followed by LiOH (0.14 mmol) and NaI (0.10 mmol). The reaction mixture was stirred at 90 °C for 16 h.
- 15 Solvent was removed *in vacuo* and the residue was dissolved in water (5 mL) and CH<sub>2</sub>Cl<sub>2</sub> (5 mL). The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3X5 mL) and the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>. Filtration and evaporation gave the crude product which was purified by flash chromatography (silica gel/ CH<sub>2</sub>Cl<sub>2</sub>:MeOH 9:1) to give the free base of the title compound.
- 20 To a solution of this material in CH<sub>2</sub>Cl<sub>2</sub> (0.2 mL) was added 0.14 mmol of HCl (1M in Et<sub>2</sub>O), the solvent evaporated *in vacuo* and the material thus obtained triturated with Et<sub>2</sub>O to give 34 mg of the title compound as a white slightly hygroscopic solid (70% yield).

- 25 **NMR (<sup>1</sup>H, MeOD):** δ 8.4 (s, 1H), 7.75 (d, 1H), 7.7 (dd, 1H), 7.41 (d, 1H), 6.6 (s, 1H), 4.0-3.84 (m, 2H), 3.80 (s, 3H), 3.78-3.70 (m, 4H), 3.52-3.18 (m, 6H) 2.51 (s, 3H), 2.47 (s, 3H). **MS (m/z):** 451.2 [MH]<sup>+</sup>.

**Example 2 : 7-(5-Methyl-3-isoxazolyl)-3-(2-([4-methyl-5-(tetrahydro-2H-pyran-4-yl)-4H-1,2,4-triazol-3-yl]thio)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride**




30



The title compound was prepared in analogy to the method described in Example 1 in 27 mg yield as a white slightly hygroscopic solid (61% yield) from 4-methyl-5-(tetrahydro-2H-pyran-4-yl)-4H-1,2,4-triazole-3-thiol (20 mg).

**NMR (<sup>1</sup>H, MeOD):** δ 7.75 (d, 1H), 7.7 (dd, 1H), 7.41 (d, 1H), 6.6 (s, 1H), 3.77 (s, 3H), 3.4 (m, 1H), 4.1 (m, 2H), 3.4 (m, 2H), 4.0-3.2 (m, 12 H), 2.5 (s, 3H), 1.99 (m, 4H). **MS (m/z):** 454.2 [MH]<sup>+</sup>


**Example 3 : 7-(5-Methyl-3-isoxazolyl)-3-(2-[[4-methyl-5-(2-methyl-5-quinolinyl)-4H-1,2,4-triazol-3-yl]thio]ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride**



The title compound was prepared in analogy to the method described in Example 1 in 30 mg yield as a white slightly hygroscopic solid (61% yield) from 4-methyl-5-(2-methyl-5-quinolinyl)-4H-1,2,4-triazole-3-thiol (27 mg).

**NMR (<sup>1</sup>H, MeOD):** δ 8.89 (d, 1H), 8.39 (d, 1H), 8.26 (t, 1H), 8.09 (d, 1H), 7.94 (d, 1H), 7.75 (d, 1H), 7.7 (dd, 1H), 7.41 (d, 1H), 6.6 (s, 1H), 4.0 (m, 2H), 3.66-3.23 (m, 10H), 3.81 (s, 3H), 3.01 (s, 3H), 2.5 (s, 3H). **MS (m/z):** 511.2 [MH]<sup>+</sup>.

**Example 4: 7-(5-Methyl-3-isoxazolyl)-3-(2-[[4-methyl-5-(2-methyl-6-quinolinyl)-4H-1,2,4-triazol-3-yl]thio]ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride**




The title compound was prepared in analogy to the method described in Example 1 in 9 mg yield as a white slightly hygroscopic solid (18% yield) from 4-methyl-5-(2-methyl-6-quinolinyl)-4H-1,2,4-triazole-3-thiol (27 mg).

**NMR (<sup>1</sup>H, MeOD):** δ 8.89 (d, 1H), 8.52 (d, 1H), 8.26 (dd, 1H), 8.2 (d, 1H), 7.87 (d, 1H), 7.63 (d, 1H), 7.57 (dd, 1H), 7.3 (d, 1H), 6.46 (s, 1H), 3.65, 3.73, 3.9-3.23 (s, s, bm, 15H), 3.73 (s, 3H), 3.65 (s, 3H), 3.01 (s, 3H), 2.5 (s, 3H). **MS (m/z):** 511.2 [MH]<sup>+</sup>.

Claims

1. A compound of formula (I) or a salt thereof:



5 wherein

- $R_1$  and  $R_4$  are independently selected from the group consisting of hydrogen, fluoro, chloro, bromo,  $C_{1-2}$ alkyl,  $C_1$ alkoxy, halo $C_{1-2}$ alkyl, halo $C_1$ alkoxy, hydroxy, cyano and nitro;

- $R_2$  and  $R_3$  are independently selected from the group consisting of:  
 10 halogen, hydroxy, cyano, nitro,  $C_{1-4}$ alkyl, halo $C_{1-4}$ alkyl,  $C_{3-6}$ cycloalkyl,  $C_{1-4}$ alkoxy, halo $C_{1-4}$ alkoxy,  $C_{1-4}$ alkoxy $C_{1-4}$ alkoxy,  $C_{1-4}$ alkylthio,  $C_{1-4}$ alkoxy $C_{1-4}$ alkyl,  $C_{3-6}$ cycloalkyl $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkoxycarbonyl,  $C_{1-4}$ alkoxycarbonyl $C_{1-4}$ alkyl,  $C_{1-4}$ alkylsulfonyl,  $C_{1-4}$ alkylsulfonyloxy, halo $C_{1-4}$ alkylsulfonyl, halo $C_{1-4}$ alkylsulfonyloxy,  $C_{1-4}$ alkylsulfonyl $C_{1-4}$ alkyl,  $C_{1-4}$ alkylsulfonamido,  $C_{1-4}$ alkylsulfonamido $C_{1-4}$ alkyl, heterocyclyl, aryl, aryl $C_{1-4}$ alkoxy, aryloxy, arylthio, arylmethyl, aroyl, aryloxymethyl, arylsulfonyl, aryl-  
 15 NR'- (wherein  $R'$  is hydrogen or  $C_{1-4}$ alkyl), arylsulfonyloxy, arylsulfonyl $C_{1-4}$ alkyl, arylsulfonamido, arylcarboxamido, arylsulfonamido $C_{1-4}$ alkyl, arylcarboxamido $C_{1-4}$ alkyl, aroyl $C_{1-4}$ alkyl, aryl $C_{1-4}$ alkanoyl, a group  $R_{11}CON(R_{12})(CH_2)_r$ ,  $R_{11}R_{12}NCO(CH_2)_r$  or  $R_{11}R_{12}NSO_2(CH_2)_r$  (in which  $r$  is 0, 1, 2, 3 or 4, and each of  $R_{11}$  and  $R_{12}$  is independently hydrogen or  $C_{1-4}$ alkyl, or in the groups  $R_{11}CON(R_{12})(CH_2)_r$ ,  $R_{11}R_{12}NCO(CH_2)_r$  and  $R_{11}R_{12}NSO_2(CH_2)_r$ ,  $R_{11}CONR_{12}$  or  $R_{11}R_{12}N$  together form a 4-, 5-, 6- or 7-  
 20 membered azacyclic group optionally containing one additional O, N or S atom in the azacycle and having 3-8 carbon atoms (including the carbon atoms contained in any optional substituent(s) of the azacycle)); wherein in any group containing an aryl moiety, the aryl may be substituted by one, two or three groups selected from the group consisting of halogen, hydroxy, cyano, nitro, amino,  $C_{1-4}$ alkyl, halo $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy, halo $C_{1-4}$ alkoxy,  $C_{1-4}$ alkylenedioxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkylsulfonyl, halo $C_{1-4}$ alkylsulfonyl,  $C_{1-4}$ alkylamino,  $C_{1-4}$ dialkylamino,  $R_{13}R_{14}NCO$  (in which  $R_{13}$  and  $R_{14}$  are independently hydrogen or  $C_{1-4}$ alkyl, or  $R_{13}R_{14}N$  together form a 4-, 5-, 6- or 7-membered azacyclic group optionally containing one additional O, N or S atom in the azacycle and having 3-8 carbon atoms (including the carbon atoms contained in any optional substituent(s) of the azacycle));  
 25  
 30  
 35

- A and B are independently N or CH;

- R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub> and R<sub>9</sub> are independently hydrogen or C<sub>1-4</sub>alkyl;
- R<sub>10</sub> is a group of the formula (a) or (b):



wherein:

- Z is C<sub>1-4</sub>alkyl, haloC<sub>1-4</sub>alkyl, C<sub>3-6</sub>cycloalkyl, phenyl, heterocyclyl, a 5- or 6-membered heteroaromatic ring or a 8- to 11-membered bicyclic group, any of which is optionally substituted by 1, 2, 3 or 4 substituents selected from the group consisting of: halogen, hydroxy, oxo, cyano, nitro, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, haloC<sub>1-4</sub>alkyl, haloC<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkylenedioxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkylsulfonyl, C<sub>1-4</sub>alkylsulfonyloxy, haloC<sub>1-4</sub>alkylsulfonyl, haloC<sub>1-4</sub>alkylsulfonyloxy, C<sub>1-4</sub>alkylsulfinyl, C<sub>1-4</sub>alkylthio, R<sub>17</sub>SO<sub>2</sub>N(R<sub>18</sub>)-, R<sub>17</sub>R<sub>18</sub>NSO<sub>2</sub>-, R<sub>17</sub>R<sub>18</sub>N-, R<sub>17</sub>R<sub>18</sub>NCO-, R<sub>17</sub>CONR<sub>18</sub>- and a 5- or 6-membered heteroaromatic ring which is optionally substituted by one or two C<sub>1-2</sub>alkyl, haloC<sub>1-2</sub>alkyl or R<sub>17</sub>R<sub>18</sub>N- (wherein R<sub>17</sub> and R<sub>18</sub> are independently hydrogen or C<sub>1-4</sub>alkyl, or R<sub>17</sub> and R<sub>18</sub> together form C<sub>3-6</sub>alkylene); and wherein substituents positioned *ortho* to one another may be linked to form a 5- or 6-membered ring; and
- R<sub>15</sub> and R<sub>16</sub> are independently hydrogen or C<sub>1-4</sub>alkyl and t is 1, 2, 3 or 4, or -(CR<sub>15</sub>R<sub>16</sub>)<sub>t</sub>- forms a C<sub>3-6</sub>cycloalkylene linker.


2. A compound as claimed in claim 1, wherein R<sub>3</sub> is hydrogen.

3. A compound as claimed in claim 1 or claim 2, wherein R<sub>2</sub> is C<sub>1-4</sub>alkyl, haloC<sub>1-4</sub>alkyl, halogen, C<sub>1-4</sub>alkylsulfonyl (e.g. methylsulfonyl or ethylsulfonyl), haloC<sub>1-4</sub>alkylsulfonyl (e.g. trifluoromethylsulfonyl), C<sub>1-4</sub>alkylsulfonyloxy (e.g. methylsulfonyloxy), haloC<sub>1-4</sub>alkylsulfonyloxy (e.g. trifluoromethylsulfonyloxy), R<sub>11</sub>R<sub>12</sub>NSO<sub>2</sub> (where each of R<sub>11</sub> and R<sub>12</sub> is independently hydrogen or C<sub>1-4</sub>alkyl or R<sub>11</sub>R<sub>12</sub>N together form a 4-, 5-, 6- or 7-membered azacyclic group optionally containing one additional O, N or S atom in the azacycle and having 3-8 carbon atoms, e.g. a piperidin-1-ylsulfonyl, pyrrolidin-1-ylsulfonyl or 1,4-morpholin-4-ylsulfonyl), a 5- or 6-membered heteroaromatic or a heterocyclyl, each of which is optionally substituted by one or two substituents selected from: halogen, cyano, C<sub>1-2</sub>alkyl (e.g. methyl or trifluoromethyl), C<sub>1-2</sub>alkoxy (e.g. methoxy), C<sub>1-2</sub>alkylenedioxy (e.g. methylenedioxy), C<sub>1-3</sub>alkanoyl (e.g. acetyl), C<sub>2</sub>alkanoylamino (e.g. acetylamino), haloC<sub>1</sub>alkylsulfonyl (e.g. trifluoromethylsulfonyl) and methylsulfonyl.

4. A compound as claimed in claim 3, wherein R<sub>2</sub> is bromo, cyano, hydroxy, chloro, methoxy, tert-butyl, methylsulfonyl, ethylsulfonyl, N,N-dimethylaminosulfonyl, pyrrolidin-1-

ylsulfonyl, 1,4-morpholin-4-ylsulfonyl, methylsulfonyloxy, pyrazin-2-yl, 5-methyl-oxazol-2-yl or 5-methyl-isoxazol-3-yl.

5. A compound as claimed in any of claims 1-4 wherein both R<sub>1</sub> and R<sub>4</sub> are hydrogen.
6. A compound as claimed in any of claims 1-5, wherein A and B are both nitrogen.
7. A compound as claimed in any of claims 1-6 wherein R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub> and R<sub>8</sub> are all hydrogen.
8. A compound as claimed in any of claims 1-7, wherein R<sub>9</sub> is methyl.
9. A compound as claimed in any of claims 1-8, wherein R<sub>10</sub> is a group of formula (a).
10. A compound as claimed in claim 9, wherein in formula (a), Z is phenyl, fluorophenyl, or quinolinyl, each of which is unsubstituted or substituted by one or more substituents selected from: halogen, or cyano, C<sub>1-2</sub>alkyl (e.g. methyl), haloC<sub>1-2</sub>alkyl (e.g. trifluoromethyl), C<sub>1-2</sub>alkoxy (e.g. methoxy), haloC<sub>1-4</sub>alkoxy (e.g. trifluoromethoxy), C<sub>1-2</sub>alkylenedioxy (e.g. methylenedioxy), C<sub>2-3</sub>alkanoyl (e.g. acetyl), C<sub>2</sub>alkanoylamino (e.g. acetylamino), methylsulfonyl, haloC<sub>1</sub>alkylsulfonyl (e.g. trifluoromethylsulfonyl), C<sub>1</sub>alkylsulfonyloxy (e.g. methylsulfonyloxy), C<sub>1</sub>alkylaminosulfonyl (e.g. methylaminosulfonyl), C<sub>1</sub>alkylsulfonylamino (e.g. methylsulfonylamino) and C<sub>1</sub>alkylaminocarbonyl (e.g. methylaminocarbonyl).
11. A compound as claimed in claim 1 having a formula (IA) or a salt thereof:



(IA)

wherein:

- A, B and R<sub>9</sub> are as defined in claim 1;
- X is a 5- or 6-membered heteroaromatic ring optionally substituted by 1, 2 or 3 substituents selected from the group consisting of: halogen, cyano, C<sub>1-2</sub>alkyl, fluoroC<sub>1-2</sub>alkyl, C<sub>1-2</sub>alkoxy, C<sub>1-3</sub>alkanoyl, C<sub>2</sub>alkanoylamino, fluoroC<sub>1</sub>alkylsulfonyl and methylsulfonyl; and
- Y is heterocyclyl, a 5- or 6-membered heteroaromatic ring or a 8- to 11-membered bicyclic group, any of which is optionally substituted by 1, 2, 3 or 4 substituents selected from the group consisting of: halogen, cyano, C<sub>1-2</sub>alkyl, haloC<sub>1-2</sub>alkyl, C<sub>1-2</sub>alkoxy, haloC<sub>1-2</sub>alkoxy, C<sub>1-2</sub>alkylenedioxy, C<sub>2-3</sub>alkanoyl, C<sub>2</sub>alkanoylamino,

methylsulfonyl, haloC<sub>1</sub>alkylsulfonyl, methylsulfonyloxy, methylaminosulfonyl, methylsulfonylamino and methylaminocarbonyl.


12. A compound as claimed in claim 1, which is:

- 7-(5-Methyl-3-isoxazolyl)-3-(2-[[4-methyl-1,3-oxazol-5yl)-4H-1,2,4-triazol-3-yl]thio}ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride, or
- 7-(5-Methyl-3-isoxazolyl)-3-(2-[[4-methyl-5-(tetrahydro-2H-pyran-4-yl)-4H-1,2,4-triazol-3-yl]thio}ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride, or
- 7-(5-Methyl-3-isoxazolyl)-3-(2-[[4-methyl-5-(2-methyl-5-quinoliny)-4H-1,2,4-triazol-3-yl]thio}ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride, or
- 7-(5-Methyl-3-isoxazolyl)-3-(2-[[4-methyl-5-(2-methyl-6-quinoliny)-4H-1,2,4-triazol-3-yl]thio}ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride

or a salt thereof.


13. A process for preparing a compound as defined in claim 1, which process comprises:

(a) reacting a compound of formula (II):



(II)


wherein R<sub>1</sub> to R<sub>8</sub> are as defined for formula (I) and L is a leaving group; with a compound of formula (III):



(III)


wherein A, B, R<sub>9</sub> and R<sub>10</sub> are as defined for formula (I); or

(b) for a compound of formula (I) wherein R<sub>2</sub> is aryl, reacting a compound of formula (IV):



wherein  $R_1$ ,  $R_3$  to  $R_{10}$ , A and B are as defined for formula (I) and W is halogen or a trifluoromethylsulfonyloxy group, or W is a group M selected from a boron derivative (e.g. a boronic acid function  $B(OH)_2$ ) or a metal function such as trialkylstannyl (e.g.  $SnBu_3$ ), zinc halide or magnesium halide; with a compound aryl- $W^1$ , wherein aryl is as defined for formula (I),  $W^1$  is halogen or a trifluoromethylsulfonyloxy group when W is a group M or  $W^1$  is a group M as defined above when W is halogen or a trifluoromethylsulfonyloxy group; or

(c) for a compound of formula (I) wherein  $R_2$  is aryloxy or arylthio, reacting a compound of formula (V):



wherein G is oxygen or sulfur, and  $R_1$ ,  $R_3$  to  $R_{10}$ , A and B are as defined for formula (I); with a reagent serving to introduce the aryl group;

and optionally thereafter for any of the steps (a), (b) or (c):

- removing any protecting group(s); and/or
- forming a salt; and/or
- converting one compound of formula (I) to a different compound of formula (I).

14. A method of treating a condition for which modulation of dopamine  $D_3$  receptors is beneficial, which comprises administering to a mammal (e.g. human) in need thereof an effective amount of a compound of a compound of any of claims 1-12.

15. A method as claimed in claim 14, wherein the condition is substance abuse and/or drug dependency.

16. Use of a compound as claimed in any of claims 1-12 in the manufacture of a medicament for the treatment of a condition in a mammal for which modulation of dopamine D<sub>3</sub> receptors is beneficial.

5

17. Use as claimed in claim 16, wherein the condition is substance abuse and/or drug dependency.

18. A compound as claimed in any of claims 1-12 for use in therapy.

10

19. A compound as claimed in any of claims 1-12 for use in the treatment of a condition in a mammal for which modulation of dopamine D<sub>3</sub> receptors is beneficial.

20. A compound as claimed in any of claims 1-12 for use in the treatment of substance abuse and/or drug dependency.

15

21. A pharmaceutical composition comprising a compound as claimed in any of claims 1-12 and a pharmaceutically acceptable carrier.

9



